## Practice procedure for the Coronavirus (SARS-CoV-2): **Definitions and Advice**

<ul> <li>face-to-face contact for &gt; 15 minutes; or</li> <li>sharing of a closed space for &gt; 2hours</li> <li>in the period extending from 24 hours before onset of symptoms in the confirmences.</li> </ul>	ed
• in the period extending from 24 hours before onset of symptoms in the confirme	ed
in the period extending from 24 hours before onset of symptoms in the committee	ed ——
case.	$\neg \neg$
Anyone with respiratory symptoms or unexplained fever should be tested for COVID-19. This is especially important for:	
anyone who lives or works in a high-risk setting, including healthcare facilities, a care and other residential facilities, schools, prisons, and other closed settings	ged
Aboriginal and Torres Strait Islander people	
people who are close contacts of a confirmed case or who have returned from overseas in the last 14 days	
• anyone admitted to hospital	
people who reside in areas of increased surveillance.	
Please make sure any health care or aged care workers or residents are noted or laboratory request form so their test can be prioritised.	the
Asymptomatic people do not require testing, except in special circumstances, e.g. they are recovered cases wishing to return to work in a health care, aged care facility, or other high-risk setting, or the person is part of a public health unit outbrea investigation.	k
all confirmed cases	
• all suspect cases as defined above	
isolation or home quarantine for 14 days?  • all people that have had COVID19 testing (to review when result available)	
all people who meet the close contact definition as above	
not accepting visitors.	
practice good hand hygiene and keep surfaces clean	
• wear a surgical mask when you are in the same room as another person	
• if possible, use a separate bathroom and toilet	
a patient can call the practice	
Hospital admission is not required unless clinically indicated.	
Home isolation guide is available	
is at least 70 years old; or	
• if the person identifies as being of Aboriginal or Torres Strait Islander descent—is least 50 years old; or	at
• is pregnant; or	
A vulnerable person:  • is the parent of a child aged under 12 months; or	
is being treated for a chronic health condition; or	
is immune compromised; or	
meets the current national triage protocol criteria for suspected COVID-19 infec	ion

Version: 5 May 2020



